

Campbell County Environmental Education Center Newsletter

Fall 2023

Bacterial Leaf Scorch of Shade Trees

Nicole Gauthier, Plant Pathology Extension Specialist

IMPORTANCE

Bacterial leaf scorch is a chronic, eventually fatal disease that has devastated many shade trees in Kentucky landscapes and street plantings since the 1980s. Since then, an increasingly large number of urban plantings and historic trees have been destroyed as this disease continues to spread.

HOSTS

The bacterial leaf scorch pathogen can infect over 100 plant species, and this list continues to grow as additional hosts and alternative hosts are identified. The most common shade tree hosts in the Commonwealth are various species of oak, maple, and sycamore. Other hosts identified in Kentucky include ash, boxelder, catalpa, dogwood, elm, gingko, hackberry, lilac, mulberry, sweetgum, and yellowwood (Table 1). In addition to these hosts, various small fruit (blackberry, blueberry, grape), herbaceous ornamentals, and weeds are known to be susceptible. Some common alternative hosts include bermudagrass, English ivy, goldenrod, Virginia creeper, and wild strawberry. **SYMPTOMS**

Infected trees may leaf-out normally in spring, but by late summer or autumn, affected leaves develop a marginal (Figure 1) or interveinal scorch (Figure 2). Depending on host, foliar scorch symptoms may appear brown, reddish-brown, or olive green (Figures 3A&B). In some hosts, a yellow or reddish-brown band may be present along the interior edge of the scorched tissue (Figures 4A,B,C,D). Affected leaves may drop prematurely or remain attached to branches. Epicormic shoots (water sprouts) can develop at the tree base or along the trunk of severely diseased trees. Affected branches with scorch symptoms may initially be randomly distributed throughout the tree, or flagging may develop on one or a few branches. As more and more branches are affected, the disease takes its toll

and causes twigs and branches to die back (Figure 5) until eventually the entire tree dies. Tree decline can seem to occur relatively guickl



Figure 1. Red oak with scorch along the leaf margin.

seem to occur relatively quickly, especially under additional environmental stresses, such as drought. However, the disease typically progresses slowly over several years. Foliar symptoms alone are not definitive for this disease since leaf scorch can also occur



Figure 2. Sycamore leaves with marginal scorch extending to interveinal leaf regions.

when trees are affected by abiotic factors (e.g. drought, transplant shock), as well as other diseases (e.g. vascular wilts, root decay). A positive diagnosis can only be determined with a laboratory analysis.

TESTING FOR BACTERIAL LEAF SCORCH

When bacterial leaf scorch is suspected, leaf tissue and petioles can be tested for the presence of the bacterium in a laboratory assay. Samples should be submitted to the UK Plant Disease Diagnostic Lab by going through a local Cooperative Extension Service office. The best time to collect samples is late summer or early autumn when the number of bacterial cells in the tissue is highest. Collect and submit at least 20 symptomatic leaves (with intact petioles) that show clear marginal scorch symptoms. In addition, it is advisable to submit branches so they can be examined for other





Figure 3. Foliar symptoms of bacterial scorch can vary greatly depending on the host. Sweetgum leaves (A) have a marginal scorch with reddish brown banding on the inner edge of the necrotic region while sugar maple (B) shows wide gray banding with a dark margin.



vascular wilt diseases, cankers, injuries, etc., in case the test for bacterial leaf scorch is negative or there are additional problems present. Samples should be submitted to an Extension office the same day they are collected.

CAUSE & DISEASE DEVELOPMENT

Bacterial leaf scorch is caused by the bacterium, Xylella fastidiosa. There are several strains or subspecies of this organism, which vary in host specificity and ability to cause disease.

Xylella fastidiosa is spread by certain sap-sucking insects in the Cicadellinae and Cercopidae families, such as leafhoppers, treehoppers, and spittlebugs.

These insects have piercing-sucking mouthparts that enable them to extract nutrients from a tree's xylem (water conducting tissue). During feeding, the insect vector either acquires the bacterium to transmit to other susceptible hosts or deposits the bacterium it has already acquired. Because the bacterium can multiply within its esophagus, the insect can continue to transmit the pathogen until it molts or dies.

Once X. fastidiosa enters a tree, it may remain undetected for several years before symptoms are apparent. Bacteria in active infections multiply in the xylem and spread upward toward branch tips. Scorch is a typical symptom of water stress and, in this case, results from the build-up of bacterial cells and host defense compounds clogging xylem vessels and slowing water uptake to leaves. Timing of symptom development in mid- to late summer in urban trees is often associated with moisture and heat stresses occurring during the growing season, placing additional stress on the host. While spread from tree-to-tree does not appear to occur very rapidly, X. fastidiosa may also be present in many other hosts, such as shrubs, grasses, and weeds growing in landscapes. Thus, the insect vectors may not necessarily pass the disease from tree-to-tree, but may initially acquire the bacterium from these alternative hosts.

DISEASE MANAGEMENT

There is no cure for bacterial leaf scorch, and trees will continue to decline over a period of several years. However, following cultural





Figure 5. Pin oak with advanced dieback symtoms.

practices that promote tree vigor may help slow the rate of decline.

Cultural Practices to Promote Tree Vigor *Prune*

Remove and destroy branches as they become infected and exhibit symptoms. It is unknown if this practice will slow disease progress, but it can help improve the tree's aesthetic value and results in the removal of hazardous limbs. *Irrigate*

Apply supplemental water during hot, dry weather to relieve water stress, which can contribute to decline.

Apply mulch

Adding a layer of mulch to the entire root zone can help preserve soil moisture and reduce water stress.

Fertilize

Apply fertilizer and adjust soil pH according to soil test results to help promote tree health. Avoid overfertilization.

Other Options

Tree replacement

Begin planting one or more non-susceptible replacement trees in the vicinity so that they will attain a reasonable size before the diseased tree must be removed. Choose native trees or those with a history of doing well in Kentucky landscapes.

Tree removal

As trees decline and become a safety hazard, complete tree removal will be necessary. *Tree injections*

While there are commercial antibiotic tree-injection products available for managing bacterial leaf scorch, these treatments lack positive long-term research results. In addition,

Common Name	Scientific Name
Blackberry	Rubus sp.
Blueberry	Vaccinium sp.
Boxelder	Acer negundo
Buckeye	Aesculus sp.
Catalpa	Catalpa speciosa
Dogwood, flowering	Cornus florida
Dogwood, oriental	Cornus kousa
Elm, American	Ulmus americana
Elm, Dutch	Ulmus x hollandica
Elm, Siberian	Ulmus pumila
Elm, wych	Ulmus glabra
Gingko	Gingko biloba
Grape, cultivated & wild	Vitis sp.
Hackberry	Celtis occidentalis
Maple, red	Acer rubrum
Maple, silver	Acer saccharinum
Maple, sugar	Acer saccharum
Lilac	Syringa vulgaris
Linden	Tilia sp.
London plane	Platanus x acerifolia
Mulberry	Morus alba
Oak, black	Quercus velutina
Oak, bluejack	Quercus incana
Oak, bur	Quercus macrocarpa
Oak, chestnut	Quercus prinus
Oak, English	Quercus robur
Oak, laurel	Quercus laurifolia
Oak, live	Quercus virginiana
Oak, northern red	Quercus rubra
Oak, pin	Quercus palustris
Oak, post	Quercus stellat
Oak, scarlet	Quercus coccinea
Oak, shingle	Quercus imbricaria
Oak, Shumard	Quercus shumardii
Oak, southern red	Quercus falcate
Oak, swamp chestnut	Quercus michauxii
Oak, swamp chestnut Oak, swamp white	Quercus bicolor
Oak, swamp white Oak, turkey	Quercus bicolor Quercus laevis
Oak, water	Quercus nigra
Oak, white	Quercus alba
Oak, willow	Quercus phellos .
Sweetgum	Liquidambar styraciflua
Sycamore	Platanus occidentalis
Yellowwood	Cladrastis kentukea

the repeated injections necessary

to slow disease progress provide ports of entry for decay organisms, which infect trees and cause additional damage. Use of tree injections is not recommended because they are not long-lasting, are expensive, and inherently damage trees.

Insecticides

Attempts to manage the insect vectors using insecticides have proven ineffective in preventing the spread of the bacterial leaf scorch pathogen.

What's Up Next?

October 18th - 10:00am Animal Adaptations October 18th - 5:30pm 4-H Nature Club November 2nd - 10:00am Winter Birding November 10th - 2:00pm Winter Birding November 15th - 2:00pm Winter Birding December 7th - 6:00pm Night Hike EVERY TUESDAY 3-4pm & FRIDAY 9-10am 1000 Hours Outside

NEW FOR 2023!

Would you like to receive our Horticulture Newsletter? Scan the QR code or call (859) 572-2600 to be added to the distribution list







Animal Adaptations

Come enjoy an indoor program on the animals of Campbell County and the unique traits that help them survive in the wilderness!

Campbell County Environmental Education Center

1261 Race Track Road Alexandria, KY 41001

October 5, 2023 @ 2:00pm ~ October 18, 2023 @ 10:00am



Registration is required. Please call (859) 694-1666 / (859) 572-2600 or register online at http://campbell.ca.uky.edu/

Cooperative Extension Service

Agriculture and Natural Resources Family and Consumer Sciences 4-H Youth Development Community and Economic Development

MARTIN-GATTON COLLEGE OF AGRICULTURE, FOOD AND ENVIRONMENT

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity. Reasonable accommodation of disability may be available with prior notice. Program information may be made available in languages other than English. University of Kentucky, Kentucky State University, U.S. Department of Agriculture, and Kentucky Counties, Cooperating. Lexington, KY 40506






Agriculture and Natural Resources Family and Consumer Sciences 4-H Youth Development Community and Economic Developme and will not discriminate on the basis of race, color ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity. Reasonable accommodation of disability, may be available with prior notice. Program information may be made available in languages other than English. University of Kentucky, Kentucky State University, U.S. Department of Agriculture, and Kentucky Counties, Cooperating, Lexington, KY 40506



Disabilities accommodated



Agriculture and Natural Resources Family and Consumer Sciences 4-H Youth Development Community and Economic Development Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity. Reasonable accommodation of disability may be available with prior notice. Program information may be made available in languages other than English. University of Kentucky, Kentucky State University, U.S. Department of Agriculture, and Kentucky Counties, Cooperating, University VA0506.



1000 HOURS OUTSIDE

@ THE CAMPBELL COUNTY ENVIRONMENTAL EDUCATION CENTER

Get in your 1000 hours outside this year with a guided walk along the nature trails!

EVERY TUESDAY 3-4PM & FRIDAY 9-10AM JANUARY - DECEMBER 2023

Some dates may be cancelled due to school field trips or inclement weather throughout the year



PLEASE CALL (859) 694-1666 / (859) 572-2600 OR REGISTER ONLINE AT HTTPS://CAMPBELL.CA.UKY.EDU/

Cooperative Extension Service

Agriculture and Natural Resources Family and Consumer Sciences 4-H Youth Development Community and Economic Development MARTIN-GATTON COLLEGE OF AGRICULTURE, FOOD AND ENVIRONMENT

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity. Reasonable accommodation of disability may be available with prior notice. Program information may be made available in languages other than English. University of Kentucky, State University, U.S. Department of Agriculture, and Kentucky Counties, Cooperating. Levinetom KY 40506



CAMPBELL COUNT FARMERS MARKET 2023

Large variety of home grown produce, breads, honey and other **KY** Proud commodities. For more information, call 859-572-2600.

Highland Heights*— Tuesdays

Senior Citizens Activity Center 3504 Alexandria Pike May 16 thru October 24 3:00 p.m. to 6:00 p.m.

Fort Thomas**— Wednesdays

Mess Hall in Tower Park 801 Cochran Avenue April 12 thru December 13 (no market on 11/22/2023) 3:00 p.m. to 6:00 p.m. Hours extend to 7:00 p.m. June-September (Senior shopping begins at 2:45 p.m.)

Alexandria*— Fridays

Southern Lanes Sports Center 7634 Alexandria Pike May 19 thru October 27 3:00 p.m. to 6:00 p.m.

Newport*— Saturdays

Next to Pepper Pod Restaurant 709 Monmouth Street May 20 thru October 28 9:00 a.m. to 12 noon

- Accepts WIC, SNAP and Senior Farmer's Market Nutrition Program
- Accepts SNAP only Supplemental Nutrition Assistance Program

Cooperative Extension Service culture and Natural Resources ily and Consumer Sciences Youth Develop nent

Educational programs of Kentucky Cooperative Extension serve all people regardless of race color, age, sex, religion, disability, or national origin. University of Kentucky, Kentucky State University, U.S. Department of Agriculture, and Kentucky Counties, Cooperating,

rou

LEXINGTON, KY 40546



